

Cosmological Constant Manipulation in Extra Dimensions for Exotic Field Propulsion

Jeff Nyman

(Original Dated: 15 November 2009)

(Initial Review: 20 January 2011)

(Revision: 30 January 2011)

(Peer Review: 28 March 2011)

(Revision: 3 April 2011)

(Candidate Acceptance: 10 May 2011)

Abstract

In this paper, I propose a new approach to generating a warp bubble metric necessary for an Alcubierre-like warp drive effect. The warp bubble would theoretically allow a spacecraft to travel at arbitrarily high velocities. Key to this idea is the ability to locally manipulate an extra dimension. String theory suggests that dimensions are globally held compact by strings wrapping around them which means it may be possible to locally increase or decrease the string tension, or even counter the effects of some string winding modes. This would change the size of the extra dimensions, allowing for faster-than-light propulsion in global reference frames. Calculations of the energy requirements of a hypothetical warp drive are put forth and a “top” speed limit is proposed based on the size of the extra dimensions.

1. Introduction

A 1994 paper written by Miguel Alcubierre [2] demonstrated that, within the framework of general relativity, a modification of space-time could be created that would allow a spacecraft to travel with arbitrarily large speeds. While the immediate idea seemed to be based on the familiar “warp drive” of *Star Trek* or the “hyperdrive” of *Star Wars*, the physical effect the paper described was more in line with the physics describing the inflationary stage of the universe. As such, the spacecraft would have a relative speed, defined as change of proper spatial distance over proper spatial time, faster than the speed of light.

Since the original paper, numerous authors have investigated and built on the original work of Alcubierre. In fact, warp drives have not been the sole interest of theoretical physicists. This was demonstrated by the formation of the [NASA Breakthrough Propulsion Program](#), the purpose of which is to investigate and expand on any ideas regarding exotic field propulsion.

The notion of a physically modelable (if not realizable) “warp drive” provides an opportunity to ask the question “what constraints do the laws of physics place on the abilities of an arbitrarily advanced civilization” [22]. In this paper I will propose an original mechanism to generate the necessary ‘warp bubble’ that all such models demand. The main focus of the paper is to demonstrate that the manipulation of the radius of one, or more, of the extra dimensions found in higher dimensional quantum gravity theories, especially those that are based on or inspired by string/M-theory, creates a local asymmetry in the cosmological constant which could be used to propel an object through space.

I find it very challenging to make predictions on how this warp drive might actually function. My route was to envision a spacecraft with an exotic power generator that could create the necessary energies to locally manipulate the extra dimension(s). In this way, an advanced spacecraft would expand/contract the compactified space-time around it, thereby creating the propulsion effect. That being said, my goal in this paper is to work on realistic model rather than a physically realizable device.

The first four sections of this paper review the necessary physics required to appreciate the new warp drive model. The remainder of the paper will introduce the propulsion concept. Calculations regarding speed limits and energy requirements will also be presented.

2. The Cosmological Constant and Quantum Field Theory

Current observations of distant supernova [13] offer further indications that the universe is expanding. This expansion is realized in Einstein's equations through the cosmological constant term, Λ .

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = 8\pi GT_{\mu\nu} - \Lambda g_{\mu\nu}$$

Equation 1

This Lambda term provides a necessary addition to the gravitational field to correlate with what is observed in nature. The fundamental origin of Lambda is still a mystery nearly a century after its introduction into cosmology. Physicists are not certain what it is that generates the cosmological constant term in Einstein's equations; we simply know that it does seem to be there. Several ideas exist as to the nature of this field. Dark energy is one very popular contemporary phrase for the Lambda term. Efforts have been made to explain the cosmological constant using the more modern quantum field theory, created after Einstein put his gravitational equations into final form [7].

Quantum field theory (QFT) is widely regarded as one of the most successful physical theories of all time. Its predictions have routinely been verified in particle accelerators and, to date, no experiment has ever shown a result that contradicts QFT. Given this, it seems only natural to try to account for the cosmological constant using QFT. The calculations [5] are based on summing all the zero point oscillations with a Planck scale cut-off which gives an estimate to the overall vacuum energy density, E_{vac} , as such:

$$\langle E_{vac} \rangle = \frac{1}{2} \int_0^M \frac{d^3k}{(2\pi)^3} \sqrt{k^2 + m^2} \approx \frac{M^4}{16\pi^2}$$

Equation 2

The energy density predicted using this equation is of order $10^{71}(\text{GeV})^4$, which conflicts with the observed value of $10^{-48}(\text{GeV})^4$. The value here is often viewed as the worst predication of theoretical physics given that it is off by a factor of 10^{119} . This failure of quantum theory has recently been re-examined using brane world scenarios that have come out of recent formulations of string theory [11], [5], [14], [1], [10], [24].

One (partial) fix to the vacuum energy calculations is the introduction of supersymmetry (SUSY). The basic idea is that all known particles have an associated "super-particle" whose spin differs by exactly one half (in \hbar units). In the case of unbroken SUSY -- when particle and super-symmetric partner have exactly equal masses -- the super-particle additions to the vacuum energy perfectly cancel the particle contributions and the resulting vacuum energy is reduced to zero.

However, when SUSY is broken and the difference of the particle mass and the super-symmetric partner mass is on a scale of $M_s \approx 10^4 \text{GeV}$, then the resulting vacuum energy is on the order $(M_s)^4 \approx 10^{16} \text{GeV}$. In

this case, the discrepancy is off by a factor of 10^{60} . That is still very large but certainly much better than the previous discrepancy. In this paper I assume that even the SUSY-breaking contribution is somehow cancelled and that all contributions to the vacuum energy density are higher dimensional.

3. Kaluza Klein Models

Kaluza-Klein (KK) theory was an early attempt to unify gravitation with electromagnetism. Theodor Kaluza initially postulated that a fifth spatial dimension could be introduced into Einstein's equations [16]. When the equations were solved the additional dimension would generate not only the standard metric, but also a vector field A_μ that could be associated with the electromagnetic field and a scalar field ϕ . Kaluza's starting point was the following metric:

$$\tilde{g}_{\tilde{\mu}\tilde{\nu}} = \begin{pmatrix} g_{\mu\nu} - \phi A_\mu A_\nu & -\phi A_\mu \\ -\phi A_\nu & -\phi \end{pmatrix}$$

Equation 3

In this metric, μ and ν run over 0,1,2,3 and the tilde quantities run over 0,1,2,3,5. A_μ and A_ν are vector fields and ϕ is a scalar field. With this formulation, you model a source free space-time (i.e., pure-gravity) in five dimensions. The corresponding action of the system is given by:

$$S^5 = - \int d^5x \sqrt{\tilde{g}} \tilde{R}$$

Equation 4

Where \tilde{g} is the five dimensional determinant of the metric and \tilde{R} is the five dimensional Ricci scalar. After integrating out the extra dimension and performing a conformal rescaling of the metric $g_{\mu\nu} \rightarrow \sqrt{\phi} g_{\mu\nu}$, it is trivial to show that the four dimensional action can be written as:

$$S = \int d^4x \sqrt{g} \left(-R + \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{4} e^{-\sqrt{3}\phi} F_{\mu\nu} F^{\mu\nu} \right)$$

Equation 5

From this action the model retrieves the spin-2 graviton, the spin-1 photon and discovers a spin-0 scalar field. This indeed was a success in unifying gravity with electromagnetism. However the unexpected existence of the scalar field was an embarrassment at the time. This stimulated later work on Brans-Dicke theories of gravity where the scalar field has the physical effect of changing the effective gravitational constant from place to place.

Kaluza's idea suffered from a very obvious drawback. If there is a fifth dimension, where is it? In 1926 Oskar Klein suggested that the fifth dimension compactifies so as to have the geometry of a circle of

extremely small radius [18]. Thus, the space has topology $R^4 \times S^1$. One way to envisage this space-time is to imagine a garden hose pipe. From a long distance it looks like a one dimensional line but a closer inspection reveals that every point on the line is in fact a circle.

4. Extra Dimensions and the Casimir Effect

The Casimir effect is one of the most salient manifestations of the vacuum fluctuations. In its most elementary form, it is the interaction of a pair of neutral, parallel conducting planes whose existence modifies the ground state of the quantum vacuum in the interior portion of the plates creating a force which attracts the plates to each other. For the electromagnetic field, this force of attraction is:

$$F = -\frac{\pi A hc}{480 a^4}$$

Equation 6

Here A is the area of the plates, h is Planck's constant, c is the speed of light and a is the plate separation. For a review on the Casimir effect see [21]. The Casimir effect can be extended to regions of non-trivial topology. For example, on S^1 -- a circular manifold -- it is possible to associate 0 and 2π with the location of the plates and the Casimir energy can be calculated. This becomes relevant when you consider models with additional spatial dimensions.

Two models with extra dimensions have become particularly popular in recent years. These are the Arkani-Hamed-Dimopoulos-Dvali (ADD) [3], [4] and the Randall-Sundrum (RS) models [22]. Both are attempts to explain what has become known as a hierarchy problem in physics, which questions why the gravitational force is so much weaker than the other forces of nature.

In the ADD model it is proposed that the force carriers of the standard model (the photon, W^+ , W^- , Z^0 and the gluons) are constrained to exist on the usual four dimensional space-time, which is called the bulk. Gravity however, is free to move both in the bulk and in the extra dimensions. These dimensions can be as large as μm [17]. Because only gravity can propagate in the extra dimensions, it is not possible to 'see' these dimensions as we see in the bulk through the electromagnetic field (light), nor can we observe them through weak or strong force interactions, which are also restricted to the bulk. It is the freedom of the graviton to propagate off of the bulk that dilutes the field strength accounting for the apparent weakness of gravity.

In the RS models, the proposal is that the warping of the extra dimension is the root cause of the weakness of gravity.

In both the ADD and RS models you can account for the cosmological constant by calculating the contribution of extra dimensional graviton fields to the Casimir energy and then associating this energy with the cosmological constant [11]. As an example I will demonstrate a calculation of the vacuum energy due to compactification in the ADD scenario in which the space-time orbifold will be $M^4 \times T^2$.

Matter fields will reside on the 3-brane (our universe) and gravity is free to propagate in the bulk. The calculations closely follow [15].

The four dimensional vacuum energy is given by:

$$\langle E_{vac} \rangle = \sum_{n_1, n_2=1}^{\infty} \frac{1}{2} \int \frac{d^3 k}{(2\pi)^3} \sqrt{k^2 + \frac{4\pi^2(n_1^2 + n_2^2)}{R^2}}$$

Equation 7

Where R is the radius of the extra dimensions and n_1 and n_2 are the KK (Kaluza Klein) graviton modes. I will now perform dimensional regularization on the k integral by first rewriting it as:

$$I(m_n) = \mu^{2\epsilon} \int \frac{d^3 k}{(2\pi)^3} (k^2 + m_n^2)^{\frac{1}{2}-\epsilon}$$

Equation 8

Here I have made the following substitution:

$$m_n^2 = \frac{4\pi^2(n_1^2 + n_2^2)}{R^2}$$

The μ term is inserted for dimensional consistency and ϵ will be taken to zero. The integral can be expressed as a beta function which can be expanded to:

$$I(m_n) = -\frac{m_n^4}{32\pi^2} \left(\frac{1}{\epsilon} + 2\log 2 - \frac{1}{2} - \log\left(\frac{m_n^2}{\mu^2}\right) \right)$$

Equation 9

The vacuum energy can thus be expressed as:

$$\langle E_{vac} \rangle = -\frac{1}{32\pi^2} \sum_{n_1, n_2=1}^{\infty} m_n^4 \left(\frac{1}{\epsilon} + 2\log 2 - \frac{1}{2} - \log\left(\frac{m_n^2}{\mu^2}\right) \right)$$

Equation 10

The summation is clearly divergent so I next use a zeta function regularization.

$$\sum_{n_1, n_2=1}^{\infty} m_n^4 = \left(\frac{4\pi}{R^2} \right)^2 \sum_{n_1, n_2=1}^{\infty} (n_1^4 + n_2^4 + n_1^2 n_2^2)$$

Equation 11

$$= \left(\frac{4\pi}{R^2} \right)^2 2 [\zeta(0)\zeta(-4) + \zeta^2(-2)]$$

Equation 12

Using the fact that $\zeta(-2m) = 0$, for m any natural number, I obtain the vacuum energy:

$$\langle E_{vac} \rangle_{4+2} = \frac{1}{32\pi^2} \sum_{n_1, n_2=1}^{\infty} m_n^4 \log\left(\frac{m_n^2}{\mu^2}\right)$$

Equation 13

$$= \frac{\pi}{2R^4} \sum_{n_1, n_2=1}^{\infty} (n_1^2 + n_2^2)^2 \log(n_1^2 + n_2^2)$$

Equation 14

$$= -\frac{\pi^2}{R^4} \zeta(0)\zeta'(4)$$

Equation 15

This result can be generalized to $4+n$ dimensions

$$\langle E_{vac} \rangle = -\frac{\pi^2}{R^4} \left[\frac{(2+n)(3+n)}{2} - 1 \right] [\zeta(0)]^{n-1} \zeta'(4)$$

Equation 16

Here the degrees of freedom of the graviton are expressed in the brackets. This vacuum energy due to the massive KK modes is associated with the cosmological constant.

5. Warp Drives

Numerous papers discussing the idea of warp drives have emerged in the literature in recent years. See for example [19]. The basic idea of all of these is to formulate a solution to Einstein's equations whereby a warp bubble is driven by a local expansion of space-time behind the bubble and a contraction ahead of the bubble. One common feature of these papers is that their physical foundation is the General Theory of Relativity. An element missing from all the papers is that there is little or no suggestion as to *how* such a warp bubble may be created.

I do not plan to buck that trend too much in that the aim of this paper is not to discuss the plausibility of a warp drive. This means I am not addressing the valid questions associated with violation of the null

energy condition or issues regarding causality. The aim of this paper is to suggest that a warp bubble could be generated using ideas and mathematics that are consistent with quantum field theory. This may have the effect of allowing us to hypothesize how such a bubble could be created by a sufficiently advanced technology.

By associating the cosmological constant with the Casimir Energy due to the Kaluza Klein modes of gravitons in higher dimensions, especially in the context of M-theory derived or inspired models, it is possible to form a relationship between Λ and the radius of the compact extra dimension. As I have shown with equation (16), the following holds:

$$\langle E_{vac} \rangle = \Lambda \propto \frac{1}{R^4}$$

Equation 17

An easier way of developing this relationship is to put things in terms of Hubble's constant H which describes the rate of expansion of space per unit distance of space.

$$H \propto \sqrt{\Lambda}$$

Equation 18

I can also put this in terms of the radius of the extra dimension, which gives:

$$H \propto \frac{1}{R^2}$$

Equation 19

This result indicates that a sufficiently advanced technology with the ability to *locally* increase or decrease the radius of the extra dimension would be able to locally adjust the expansion and contraction of space-time creating the hypothetical warp bubble that I discussed in earlier sections. A spacecraft with the ability to create such a bubble will always move inside its own local light-cone. However the ship can utilize the expansion of space-time behind the ship to move away from some object at any desired speed or equivalently to contract the space-time in front of the ship to approach any object.

In the context of general relativity a similar phenomenology is produced for the case of anisotropic cosmological models, in which it is the contraction of the extra dimension that has the effect of expanding another [19]. A common example is to consider a 'toy' universe with one additional spatial dimension with the following metric:

$$ds^2 = dt^2 - a^2(t)d\vec{x}^2 - b^2(t)dy^2$$

Equation 20

In this toy universe you can assume space-time is empty, that there is no cosmological constant and that all spatial dimensions are locally flat. The metric is thus:

$$T_{\mu\nu} = \Lambda g_{\mu\nu} = 0$$

Equation 21

The action of the Einstein theory of gravity generalized to five dimensions will be

$$S^{(5)} = \int d^4x dy \sqrt{-g^{(5)}} \left(\frac{M_5^2}{16\pi} R^{(5)} \right)$$

Equation 22

Solving the vacuum Einstein equations gives:

$$G_{\mu\nu} = 0$$

Equation 23

I obtain for the G_{11} component as such:

$$G_{11} = \frac{3\dot{a}(b\dot{a} + a\dot{b})}{a^2b}$$

Equation 24

Rewriting $\dot{a}/a = H_a$ and $\dot{b}/b = H_b$ where H_a and H_b corresponds to the Hubble constant in three space and the Hubble constant in the extra dimension respectively, I find that solving for $G_{11} = 0$ yields:

$$H_a = -H_b$$

Equation 25

This remarkable result indicates that in a vacuum, the shear of a contracting dimension is able to inflate the remaining dimensions. In other words the expansion of the 3-volume is associated with the contraction of the one-volume.

Even in the limit of flat space-time with zero cosmological constant, general relativity shows that the physics of the compactified space affects the expansion rate of the non-compact space. The main difference to note here is that the quantum field theoretic result demonstrates that a fixed compactification radius can also result in expansion of the three-volume as is shown in equation (16)

due to the Casimir effect, whereas the general relativistic approach suggests that a changing compactification radius results in expansion. Both add credibility to the warp drive concept presented in this paper.

6. Energy Requirements

In this section I will perform some elementary calculations to determine how much energy would be required to reach superluminal speeds. I will also determine an absolute speed limit based on fundamental physical limitations.

The currently accepted value for the Hubble constant is 70 km/sec/Mpsc. A straightforward conversion into SI units gives $H = 2.17 \times 10^{-18} \text{ (m/s)/m}$. This tells us that one meter of space expands to two meters of space if one were prepared to wait two billion billion seconds or sixty five billion years. The fundamental idea behind the warp drive presented in this paper is to increase Hubble's constant such that space no longer expands at such a slow rate, but locally expands at an arbitrarily fast velocity. For example, if I want space to locally expand at the speed of light, a simple calculation shows by what factor I would need to increase H.

$$\frac{H_c}{H} \approx \frac{10^8}{10^{-18}} = 10^{26}$$

Equation 26

Here H_c is the 'modified' Hubble constant (where c is the speed of light). This result implies that H would have to be increased by a factor of 10^{26} for space to expand at the speed of light. Since it is known that $H \propto 1/R^2$, it is possible to (perhaps naively) form the relation

$$\frac{H_c}{H} = \frac{R^2}{R_c^2} = 10^{26}$$

Equation 27

Or I could write this as solving for the radius:

$$R_c = 10^{-13} R$$

Equation 28

Here R_c is the modified radius of the extra dimension. This indicates that the extra dimensional radius must be locally reduced by a factor of 10^{13} to stimulate space to expand at the speed of light. In the ADD model the size of the extra dimension can be as large as 10^{-6} m . If I use this number as a prototype extra-dimensional radius, this would have to be shrunk to 10^{-19} m for light speed expansion.

I can now look at the energy required to create the necessary warp bubble. The accepted value of the cosmological constant is $\Lambda \approx 10^{-47}(\text{GeV})^4$. Converting again into SI units gives $\Lambda \approx 10^{-10}\text{J}/\text{m}^3$. Now, for a warp bubble expanding at the speed of light I would need to increase this again by a factor of 10^{52} as I have $H \propto \sqrt{\Lambda}$. I can thus say this:

$$\Lambda_c = 10^{52}\Lambda = 10^{42}\text{J}/\text{m}^3$$

Equation 29

Here Λ_c is the local value of the cosmological constant when space is expanding at c . To make this a concrete example, I will consider a spacecraft of these dimensions:

$$V_{craft} = 10\text{m} \times 10\text{m} \times 10\text{m} = 1000\text{m}^3$$

Equation 30

If I postulate that the warp bubble must, at least, encompass the volume of the craft, the total amount of energy ‘injected’ locally would equal

$$E_c = \Lambda_c \times V_{craft} = 10^{45}\text{J}$$

Equation 31

Assuming some arbitrarily advanced civilization was able to create such an effect, I will further postulate that this civilization would be able to utilize the most efficient method of energy production, which would be matter/antimatter annihilation. Using $E = mc^2$ this warp bubble would require around 10^{28}Kg of antimatter to generate, roughly the mass-energy of the planet Jupiter.

This energy requirement would drop dramatically if I assumed a thin-shell of modified space-time instead of bubble encompassing the volume of the craft. What this shows is that larger ships would be more problematic than smaller vessels.

7. Speed Limits

It is known from string theory that the absolute minimum size for an extra dimension is the Planck length, 10^{-35}m . This will place an ultimate speed limit on the expansion of space based on the idea that there is a limit to the minimum radius of the extra dimension. From the above arguments it is straightforward to form the relation

$$\frac{H_{max}}{H} = \frac{R^2}{R_{min}^2} = \frac{10^{-12}}{10^{-70}} = 10^{58}$$

Equation 32

Here H_{max} is the maximum rate of space-time expansion based on the minimum radius of the extra dimension R_{min} . In this formula I have again used a prototype extra-dimensional radius of 10^{-6} m which is the upper bound based on current experimental limits. Using these values and the known value of H in SI units I obtain the following expansion value:

$$H_{max} = 10^{58} H \approx 10^{40} (m/s)/m$$

Equation 33

A quick conversion into multiples of the speed of light reveals this velocity:

$$V_{max} = 10^{32} c$$

Equation 34

Such a velocity would require on the order of 10^{99} Kg of antimatter. On the negative side, this is more mass energy than is contained within the universe. On the positive side, at this velocity it would be possible to cross the known universe in a little over 10^{-15} seconds.

I do want to reiterate that the calculations in the previous two sections are extremely 'back of the envelope' and merely serve as interesting figures to contemplate, based on the vacuum energy given by:

$$\langle E_{vac} \rangle = \Lambda \propto \frac{1}{R^4}$$

Equation 35

Note that, as far as it is possible to know, it is not really possible to travel faster than light in a local sense. This means that, in accordance with relativity, faster than light restrictions do apply in local reference frames. It is possible, however, to make a round trip between two points in an arbitrarily short time as measured by an observer who remained at rest at the starting point and this is achieved by the global manipulation of space-time, outside of local reference frames. Even with that approach, others [24] have discussed the many details on violations of the null energy conditions and causality.

8. Conclusions

In this paper I have calculated the vacuum energy due to extra dimensional graviton contributions to the Casimir energy and associated this energy with the cosmological constant. It has been shown that this energy is intimately related to the size of the extra dimension. I have picked the ADD model, however similar approaches can be used for alternative models, for example the RS model of warped extra dimension where a similar relation can be found.

I have proposed that a sufficiently advanced civilization could utilize this relation to generate a localized expansion/contraction of space-time creating a 'warp bubble' in which to travel at arbitrarily high velocities. Further research is required to determine the mathematical and physical basis for how to locally manipulate an extra dimension. String theory suggests that dimensions are globally held compact by strings wrapping around them [6], [9]. If this is indeed the case, then it may be possible to even locally increase or decrease the string tension, or even locally counter the effects of some string winding modes. This would achieve the desired effect of changing the size of the extra dimensions which would lead to propulsion under this model.

Further research will also be useful to learn how to perform calculations as to the energies required to affect an extra dimension and to try and relate this energy to the acceleration a spacecraft might experience.

9. Bibliography

- [1] Aghababaie, Y., Burgess, C., Parameswaran, S., Quevedo F., "Towards a Naturally Small Cosmological Constant from Branes in 6-D Supergravity," *Nucl.Phys. B680*, 389-414, (2004).
- [2] Alcubierre, M., "The Warp Drive: Hyperfast Travel Within General Relativity," *Class.Quant.Grav.Letters 11*, L73-L77, (1994).
- [3] Antoniadis, I, Arkani-Hamed, N., Dimopoulos, S., Dvali G., "New dimensions at a millimeter to a Fermi and superstrings at a TeV". *Phys. Lett. B 436*: 257-263, (1998).
- [4] Arkani-Hamed, N., Dimopoulos, S., Dvali G., "The Hierarchy Problem and New Dimensions at a Millimeter". *Phys. Lett. B 436*: 263-272, (1998).
- [5] Binetruy, P., Deffayet, C., Ellwanger, U., Langlois, D., "Brane Cosmological Evolution in a Bulk with Cosmological Constant," *Phys.Lett. B477*, 285-291, (2000).
- [6] Brandenberger, R., and Vafa, C., "Superstrings in the Early Universe," *Nucl. Phys. B136*, 391 (1989)
- [7] Chen, P., and Gu, J., "Casimir Effect in a Supersymmetry-breaking Brane World as Dark Energy", *SLAC-PUB-10772*, 2004.
- [8] Cherednikov, I., "On Casimir Energy Contribution to Observable Value of the Cosmological Constant," *Acta Phys. Polon. B33*:1973-1977, (2002).
- [9] Cleaver, G and Rosenthal P., "String Cosmology and the Dimension of Space-time," *Nucl.Phys. B457*, 621-642, (1995).
- [10] de Alwis, S., "Brane World Scenarios and the Cosmological Constant," *Nucl. Phys. B597*, 263-278, (2001).
- [11] Ellwanger, U., "Brane Universes and the Cosmological Constant," *Mod.Phys.Lett. A20*, 2521-2532, (2005).
- [12] Garriga, J., Pujolas, O., Tanaka, T., "Radion Effective Potential in the Brane World," *Nucl.Phys. B605*, 192-214, (2001).
- [13] Garnavich, P., "Observational Evidence From Supernovae for an Accelerating Universe and a Cosmological Constant," *Astron.J. 116*:1009-1038, (1998).
- [14] Giudice, G., and Rattazzi, R., "Theories with Gauge Mediated Supersymmetry Breaking," *Phys. Rept. 322*, 419-499, (1999).
- [15] Gupta, A., "Contribution of Kaluza-Klein Modes to the Vacuum Energy in Models with Large Extra Dimensions and the Cosmological Constant", eprint arXiv:hep-th/0210069
- [16] Kaluza, T., Sitz. Preuss. Akad. Wiss. *Phys. Math. Kl 996* (1921).

- [17] Kapner, et al., "Test of the Gravitational Inverse-Square Law below the Dark-Energy Length Scale," *Phys. Rev. Lett.* **98** 021101 (2007).
- [18] Klein, O., *Z. F. Physik* **37** 895 (1926); Klein, O., *Nature* **118** 516 (1926)
- [19] Levin, J., "Inflation from Extra Dimensions," *Phys.Lett.* **B343** 69-75, (1995).
- [20] Lobo, S., Visser M., "Fundamental limitations on 'warp drive' spacetimes," *Class.Quant.Grav.***21**:5871-5892,2004.
- [21] Milton, K., "The Casimir effect: Recent controversies and progress," *J.Phys.***A37**:R209,2004
- [22] Morris, M., and Thorne, K., and Yurtsever, U., "Worm-holes, Time Machines, and the Weak Energy Conditions," *Phys. Rev. Lett.*, **1446-1449**, (1988).
- [23] Randall, L., and Sundrum, R., "An Alternative to Compactification," *Phys. Rev. Lett.* **83**, 4690 (1999);
- [24] Tseytlin, A., and Vafa, C., "Elements of String Cosmology," *Nucl. Phys.* **B372**, 443-466, (1992).
- [25] Visser, M., and Bassett, B., and Liberati, S., "Perturbative superluminal censorship and the null energy condition," arXiv:gr-qc/9908023v1 (1999).